

Glimitab-MV $\frac{1}{2}$

1. Generic Name

Glimepiride +Metformin + Voglibose

2. Quantitative and Qualitative Composition

Each tablet contains:

Glimepiride	1mg/2mg
Metformin SR	500mg
Voglibose	0.2mg

3. Dosage form and strength

Glimitab- MV is available in the strengths of 1/2mg.

4. Clinical particulars

4.1 Therapeutic indication

Glimitab MV is indicated as an adjunct to diet and exercise to improve glycaemic control in patients with type-2 diabetes who are already treated with a combination of glimepiride and metformin or whose diabetes is not adequately controlled with metformin alone or require additional glycaemic control.

4.2 Posology and method of administration

- Dosage should be individualized on the basis of both effectiveness and tolerance. The combination should be given once daily before meals and should be started at a low dose. Starting dose for patients inadequately controlled on Metformin monotherapy or Glimepiride and Metformin dual therapy.
- Glimitab MV may be initiated once daily, and gradually titrated after assessing the therapeutic response.



We Impart Health to Life

- Glimitab MV can be starting dose for patients who initially responded to glimepiride monotherapy or Glimepiride and Metformin dual therapy and require additional glycaemic control.
- Based on the initial starting dose of Glimitab MV may be initiated once daily, and gradually titrated after assessing the therapeutic response.

4.3 Contraindication

- Renal disease or renal dysfunction, as suggested by serum creatinine levels ≥ 1.5 mg/dL [males], ≥ 1.4 mg/dL [females] or abnormal creatinine clearance, which may also result from conditions such as cardiovascular collapse (shock), acute myocardial infarction, and septicaemia.
- Hepatic impairment.
- Known hypersensitivity to this product or any of its components.
- Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma. Diabetic ketoacidosis should be treated with insulin.
- Patients undergoing radiologic studies involving intravascular administration of iodinated contrast materials, because the use of such products may result in acute alteration of renal function.
- Gastrointestinal obstruction or predisposed to it

4.4 Special warnings and precautions for use

- **Cardiac effects:** -The administration of oral hypoglycaemic drugs has been reported to be associated with increased cardiovascular mortality as compared to treatment with diet alone or diet plus insulin. In view of close similarities between the oral hypoglycaemic drugs, this warning also applies for glimepiride. Palpitations, Cardiac flutter, Myocardial infarction, Cardiac failure are some of cardiac side effects due to glimepiride.
- **Lactic acidosis:** Lactic acidosis is a rare, but serious, metabolic complication that can occur due to metformin accumulation during treatment with glimepiride and metformin combination therapy; when it occurs, it is fatal in approximately 50% of cases. When metformin is implicated as the cause of lactic acidosis, metformin plasma



levels $>5 \mu\text{g/mL}$ are generally found. The reported incidence of lactic acidosis inpatients receiving metformin hydrochloride is very low (approximately 0.03cases/1000 patient-years, with approximately 0.015 fatal cases/1000 patient years). Reported cases have occurred primarily in diabetic patients with significant renal insufficiency and congestive heart failure.

- Lactic acidosis is a medical emergency that must be treated in a hospital setting. In a patient with lactic acidosis, metformin should be discontinued immediately and general supportive measures promptly instituted. Because metformin hydrochloride is dialyzable, prompt haemodialysis is recommended to correct the acidosis and remove the accumulated metformin.
- **Hypoglycaemia:** Most reported adverse effect with sulphonylurea drugs is severe hypoglycaemia. Proper patient selection, dosage, and instructions are important to avoid hypoglycaemic episodes.
- **Loss of control of blood glucose:** When a patient stabilized on any diabetic regimen, is exposed to stress such as fever, trauma, infection, or surgery, a temporary loss of glycaemic control may occur. At such times, it may be necessary to withhold the diabetic regime and temporarily administer insulin. The oral antidiabetic therapy may be reinstated after the acute episode is resolved.
- **Alcohol intake:** Alcohol is known to potentiate the effect of metformin on lactate metabolism. Patients should be warned against excessive alcohol intake, while receiving metformin.
- **Hypoxic states:** Cardiovascular collapse (shock) from whatever cause, acute congestive heartfailure, acute myocardial infarction and other conditions characterized by hypoxemia have been associated with lactic acidosis and may also cause prerenal azotaemia. When such events occur in patients on metformin therapy, the drug should be promptly discontinued.
- **Haemolytic anaemia:** Treatment of patients with glucose 6-phosphate dehydrogenase (G6PD) deficiency with sulfonylurea agents can lead to haemolytic anaemia. Since glimepiride is a sulfonylurea agent, caution should be used in patients with G6PD deficiency and anon-sulfonylurea alternative should be considered.

4.5 Drug interactions

- **Cationic drugs:** Certain medications used concomitantly with metformin may increase the risk of lactic acidosis. Cationic drugs that are eliminated by renal tubular secretions (e.g: amiloride, digoxin, morphine, procainamide, quinidine, ranitidine, or vancomycin) may decrease metformin elimination by competing for common renal tubular transport systems. Hence, careful patient monitoring and dose adjustment of metformin/cationic drug is recommended.
- **Miconazole (systemic route, oromucosal gel) and Phenylbutazone (systemic route):** Increases hypoglycaemic effect of glimepiride.
- **Furosemide:** A single-dose, metformin-furosemide drug interaction study in healthy subjects demonstrated that pharmacokinetic parameters of both compounds were affected by co-administration. Furosemide increased the metformin plasma and blood C_{max} by 22% and blood AUC by 15%, without any significant change in metformin renal clearance. When administered with metformin, the C_{max} and AUC of furosemide were 31% and 12% smaller, respectively, than when administered alone, and the terminal half-life was decreased by 32%, without any significant change in furosemide renal clearance.
- **Vitamin B12:** Metformin may result in suboptimal oral vitamin B12 absorption by competitively blocking the calcium dependent binding of the intrinsic factor vitamin B12 complex to its receptor. The reaction very rarely results in pernicious anaemia which is reversible with discontinuation of metformin and supplementation with vitamin B12.
- **Nifedipine:** Nifedipine appears to enhance the absorption of metformin, it increases plasma metformin C_{max} and AUC by 20% and 9% respectively and increases the amount of metformin excreted in the urine.
- **Danazol:** If the use of this active substance cannot be avoided, the patient's urine and blood glucose must be monitored. It may be necessary to adjust the dose of glimepiride and metformin during and after treatment with danazol.
- **Salicylates:** If salicylates are administered or discontinued in patients receiving oral antidiabetic agents, patients should be monitored for hypoglycaemia or loss of blood glucose control.



- **Thiazide:** Interactions between thiazide diuretics and oral antidiabetic agents decreases insulin sensitivity thereby leading to glucose intolerance and hyperglycaemia. Hence diabetic patients should be monitored closely.
- **Other:** Concomitant administration of angiotensin enzyme inhibitors (captopril, enalapril), other antidiabetic drugs (insulin, acarbose) beta-blockers, fluconazole, monoamine oxidase inhibitors (MAOIs), sulphonamides and NSAIDs, increases sensitivity to insulin and potentiates blood glucose lowering effect and may in some instances, cause hypoglycaemia. Patients receiving estrogens or oral contraceptives, phenytoin, quinolones should be closely monitored for loss of diabetic control.

4.6 Use in special population

- **Pediatric:** Safety and effectiveness of Glimitab-MV combination in paediatric patients have not been established
- **Geriatric:** Aging is associated with reduced renal function; Glimitab-MV should be used with caution in the elderly.
- **Liver impairment:** should be administered with extreme caution in patients with impaired liver function. It is advised to consult your doctor to make a suitable adjustment in dosage and/or time-interval between two dosages.
- **Renal failure:** Metformin is known to be excreted by the kidneys, and because risk of serious adverse reactions to the drug is greater in patients with impaired renal function, Glimitab-MV should be used only in patients with normal renal function.
- **Pregnancy and lactation:** Abnormal blood glucose levels during pregnancy are associated with the higher incidence of congenital abnormalities. Most experts suggest insulin be used to maintain the blood glucose levels as close to normal as possible. The use of glimepiride and metformin combination is not recommended for use in pregnancy.

Studies in lactating rats show that metformin is excreted into milk and reaches levels comparable to those in plasma. Similar studies have not been conducted on nursing mothers. Also Glimepiride should not be used by breast-feeding mothers. Hence, the use of glimepiride and metformin combination is not

recommended for use in lactating mothers, and if the diet alone is inadequate for controlling blood glucose, insulin therapy should be considered.

4.7 Effects on ability to drive and use machine

No data available.

4.8 Undesirable effects

- **Glimepiride:** Thrombocytopenia, Palpitations, Vertigo, Vision blurred, Nausea, Diarrhoea, Vomiting, Dyspepsia, Abdominal pain, Constipation, Abdominal discomfort, Hepatic function abnormal, Drug hypersensitivity, Hypoglycaemia, Hyperglycaemia, Decreased appetite, Diabetes mellitus inadequate control, Dizziness, Hypoglycaemic coma, Headache, Somnolence, Tremor, Loss of consciousness, Confusional state, Acute kidney injury, Dyspnoea, Pruritus, Rash, Hyperhidrosis, Urticaria, Hypertension
- **Metformin:** Anaemia, Thrombocytopenia, Leukocytosis, Palpitations, Cardiac arrest, Myocardial infarction, Tachycardia, Bradycardia, Cardio-respiratory arrest, Cardiac failure, Atrial fibrillation, Cardiac failure congestive, Cardiac disorder, Cardiac flutter, Arrhythmia, Vertigo, Tinnitus, Vision blurred, Visual impairment, Periorbital oedema, Diarrhoea, Nausea, Vomiting, Abdominal pain, Abdominal discomfort, Dyspepsia, Abdominal distension, Abdominal pain upper, Flatulence, Constipation, Gastrointestinal disorder, Gastritis, Dry mouth, Gastroesophageal reflux disease, Pancreatitis, Pancreatitis acute, Gastric disorder, Eructation, Dysphagia, Epigastric discomfort, Retching, Gastric dilatation, Gastrointestinal pain, Frequent bowel movements, Lip swelling, Abnormal faeces, Hepatic function abnormal, Hepatitis, Drug-induced liver injury, Jaundice, Liver disorder, Hepatic failure, Hepatic steatosis, Drug hypersensitivity, Hypersensitivity, Lactic acidosis, Hypoglycaemia, Decreased appetite, Metabolic acidosis, Hyperglycaemia, Diabetes mellitus inadequate control, Hyperkalaemia, Dehydration, Diabetic ketoacidosis, Diabetes mellitus, Acidosis, Vitamin B12 deficiency, Hyponatraemia, Hyperlactacidaemia, Hypokalaemia, Appetite disorder, Ketoacidosis, Fluid retention, Type 2 diabetes mellitus, Hypophagia, Myalgia,



We Impart Health to Life

Muscle spasms, Arthralgia, Pain in extremity, Back pain, Muscular weakness, Rhabdomyolysis, Joint swelling, Arthritis, Dizziness, Headache, Somnolence, Completed suicide, Confusional state, Insomnia, Anxiety, Acute kidney injury, Renal failure, Renal impairment, Dyspnoea, Cough, Pruritus, Rash, Urticaria, Hypotension, Hypertension, Shock.

- **Voglibose:** Diarrhoea, Hepatic function abnormal, Hypoglycaemia, Dizziness

4.9 Overdose

Overdosages of sulfonylureas, including glimepiride, can produce hypoglycaemia. Mild hypoglycaemic symptoms without loss of consciousness or neurologic findings should be treated aggressively with oral glucose and adjustments in drug dosage and/or meal patterns. Close monitoring should continue until the physician is assured that the patient is out of danger. Severe hypoglycaemic reactions with coma, seizure, or other neurological impairment occur infrequently, but constitute medical emergencies requiring immediate hospitalization. If hypoglycaemic coma is diagnosed or suspected, the patient should be given a rapid IV injection of concentrated (50%) glucose solution. This should be followed by a continuous infusion of a more dilute (10%) glucose solution at a rate that will maintain the blood glucose level above 100 mg/dL. Patients should be closely monitored for a minimum of 24 to 48 hours, because hypoglycaemia may recur after apparent clinical recovery. Lactic acidosis is a rare, but serious, metabolic complication that can occur if metformin accumulates during treatment due to overdosing. Strict monitoring should be continued until the doctor is sure that the patient is out of danger. Voglibose competitively and reversibly inhibits the α -glucosidase enzymes (glucoamylase, sucrase, maltase, and isomaltase) in the brush border of the small intestine, which delays the hydrolysis of complex carbohydrates. It is unlikely to produce hypoglycaemia in overdose, but abdominal discomfort and diarrhoea may occur.

5. Pharmacological properties

5.1 Mechanism of action

- The mechanism of action of glimepiride in lowering blood glucose appears to be dependent on stimulating the release of insulin from functioning pancreatic beta cells,



We Impart Health to Life

and increasing sensitivity of peripheral tissues to insulin. Glimepiride likely binds to ATP-sensitive potassium channel receptors on the pancreatic cell surface, reducing potassium conductance and causing depolarization of the membrane. Membrane depolarization stimulates calcium ion influx through voltage-sensitive calcium channels. This increase in intracellular calcium ion concentration induces the secretion of insulin.

- Metformin improves glucose tolerance in patients with type-2 diabetes (NIDDM), lowering both basal and postprandial plasma glucose. Metformin decreases hepatic glucose production, decreases intestinal absorption of glucose, and improves insulin sensitivity by increasing peripheral glucose uptake and utilization. Hence, the combination of glimepiride and metformin sustained-release complements each other and provides better glycaemic control in management of type-2 diabetes and probably in the prevention of its associated macrovascular and microvascular complications.
- Alpha-glucosidase inhibitors are saccharides that act as competitive inhibitors of enzymes needed to digest carbohydrates: specifically alpha-glucosidase enzymes in the brush border of the small intestines. The membrane-bound intestinal alpha-glucosidase hydrolyse oligosaccharides, trisaccharides, and disaccharides to glucose and other monosaccharides in the small intestine. Acarbose also blocks pancreatic alpha-amylase in addition to inhibiting membrane-bound alpha-glucosidase. Pancreatic alpha-amylase hydrolyses complex starches to oligosaccharides in the lumen of the small intestine. Inhibition of these enzyme systems reduces the rate of digestion of complex carbohydrates. Less glucose is absorbed because the carbohydrates are not broken down into glucose molecules. In diabetic patients, the short-term effect of these drugs therapies is to decrease current blood glucose levels: the long term effect is a small reduction in hemoglobin-A1c level.

5.2 Pharmacodynamics properties

- Glimepiride, like glyburide and glipizide, is a "second-generation" sulfonylurea agent. Glimepiride is used with diet to lower blood glucose by increasing the secretion of insulin from pancreas and increasing the sensitivity of peripheral tissues to insulin.

- Metformin decreases hepatic glucose production, decreases intestinal absorption of glucose, and improves insulin sensitivity by increasing peripheral glucose uptake and utilization. Unlike sulfonylureas, metformin does not produce hypoglycaemia in either patients with type 2 diabetes or normal subjects (except in special circumstances) and does not cause hyperinsulinemia. With metformin therapy, insulin secretion remains unchanged while fasting insulin levels and day-long plasma insulin response may actually decrease
- Voglibose is an alpha glucosidase inhibitor which reduces intestinal absorption of starch, dextrin, and disaccharides by inhibiting the action of α -glucosidase in the intestinal brush border. Inhibition of this enzyme catalyses the decomposition of disaccharides into monosaccharides and slows the digestion and absorption of carbohydrates; the post-prandial rise in plasma glucose is blunted in both normal and diabetic subjects resulting in improvement of post prandial hyperglycaemia and various disorders caused by hyperglycaemia. α -Glucosidase inhibitors do not stimulate insulin release and therefore do not result in hypoglycaemia. Hence, the combination of glimepiride, metformin sustained-release and Voglibose complements each other and provides better glycaemic control in management of type-2 diabetes and probably in the prevention of its associated macrovascular and microvascular complications.

5.3 Pharmacokinetic properties

- Glimepiride is completely absorbed from the gastrointestinal tract. Peak plasma concentrations occur in 2 to 3 hours, and it is highly protein bound. The drug is extensively metabolised to two main metabolites, a hydroxy derivative and a carboxy derivative. The half-life after multiple doses is about 9 hours. About 60% of a dose is eliminated in the urine and 40% in the faeces.
- Metformin hydrochloride is slowly and incompletely absorbed from the gastrointestinal tract; the absolute bioavailability of a single 500-mg dose is reported to be about 50 to 60%, although this is reduced somewhat if taken with food. Once absorbed, protein binding in plasma is negligible; the drug is excreted unchanged in the urine. The plasma elimination half-life is reported to range from about 2 to 6 hours



after oral doses. Metformin crosses the placenta and is distributed into breast milk in small amounts.

- Voglibose is poorly absorbed after oral doses. Plasma concentrations after oral doses have usually been undetectable. After an 80 mg dose (substantially higher than recommended dose), peak plasma levels of about 20 ng/mL were observed in 1 to 1.5 hours. After ingestion of Voglibose (and other glucosidase inhibitors), the majority of active unchanged drug remains in the lumen of the gastrointestinal tract to exert its pharmacological activity. Voglibose is metabolized by intestinal enzymes and by the microbial flora. Voglibose is excreted in the urine and faeces. In a study in which a single dose of 1 mg/kg of C14-Voglibose was administered to rats, the transfer of Voglibose to the foetus and to mother's milk was observed, and the rates of excretion into urine and faeces were about 5% and 98%, respectively.

6. Nonclinical properties

6.1 Animal Toxicology or Pharmacology

Not required.

7. Description

- **Glimepiride**

Glimepiride is a long-acting, third-generation sulfonylurea with hypoglycemic activity. Compared to other generations of sulfonylurea compounds, glimepiride is very potent and has a longer duration of action.

Chemical Name -

4-ethyl-3-methyl-N-[2-[4-[(4-methylcyclohexyl)carbamoylsulfamoyl]phenyl]ethyl]-5-oxo-2H-pyrrole-1-carboxamide

Molecular Weight- 490.6 g/mol

Molecular Formula- C₂₄H₃₄N₄O₅S


Structure-

Chemical Name - (1S,2S,3R,4S,5S)-5-(1,3-dihydroxypropan-2-ylamino)-1-(hydroxymethyl)cyclohexane-1,2,3,4-tetrol

Molecular Weight- 267.28 g/mol

Molecular Formula- C₁₀H₂₁NO₇

Structure-



8. Pharmaceutical particulars

8.1 Incompatibilities

There are no known incompatibilities.

8.2 Shelf-life

18 months.

8.3 Packaging Information

Glimitab- MV is available in the strip of 10 tablets.

8.4 Storage and handling instructions

Store in a cool and dry place.



We Impart Health to Life

9. Patient Counselling Information

9.1 Adverse Reactions

Refer part 4.8

9.2 Drug Interactions

Refer part 4.5

9.3 Dosage

Refer part 4.2

9.4 Storage

Refer part 8.4

9.5 Risk Factors

Refer part 4.4

9.6 Self-monitoring information

NA

9.7 Information on when to contact a health care provider or seek emergency help

Patients are advised to be alert for the emergence or worsening of the adverse reactions and contact the prescribing physician.

9.8 Contraindications

Refer part 4.3

10. Manufactured by PURE AND CURE HEALTHCARE PVT LTD

11. Details of permission or license number with date

License No. 31/UA/2013 & 51/UA/SC/P-2013 , dated 30.11.2013

12. Date of revision: January 2022



We Impart Health to Life



We Impart Health to Life