



1. Generic name: Phenylephrine Hydrochloride, Chlorpheniramine Maleate, Sodium Citrate, Menthol

2. Qualitative and Quantitative composition

Each 5 ml. (teaspoonful) contains:

Phenylephrine Hydrochloride IP 5 mg
Chlorpheniramine Maleate IP 1 mg
Sodium Citrate IP 60 mg
In mentholated syrupy base

Colour: Carmoisine

3. Dosage form and strength

Dosage form: Available in syrup form.

Strength: Syrup containing Phenylephrine Hydrochloride 5 mg, Chlorpheniramine Maleate 1 mg, Sodium Citrate 60 mg.

4. Clinical particulars

4.1 Therapeutic indication:

Sinarest AF Jr. syrup is indicated for treatment of allergic rhinitis and common cold.

4.2 Posology and method of administration

The usual recommended oral dose of Sinarest AF Jr. Syrup is:

2-5 years = 5ml Three/ Four times a day. 6-12 years

= 10ml Three/ Four times a day.

>12 years = 10 - 15ml Three/ Four times a day.

4.3 Contraindication

The use of Sinarest AF Jr. syrup is contraindicated in patients with:

- Hypersensitivity to any of the ingredients of the formulation.
- Severe hypertension.

4.4 Special warnings and precautions for use

- “The fixed dose combination shall not be used in children below four years of age”
- Caution in elderly patients, hyperthyroidism, myocardial disease, bradycardia, partial heart block or severe arteriosclerosis when administering phenylephrine.
- Caution in asthma, bladder neck obstruction, cardiovascular disease, COPD, GI obstruction, glaucoma, hepatic impairment, hyperthyroidism, increased intraocular pressure, malnutrition, renal impairment, elderly patients, and patients taking CNS depressants.
- Phenylephrine should be used with care in patients with closed angle glaucoma and prostatic enlargement.
- Chlorpheniramine, in common with other drugs having anticholinergic effects, should be used with caution in epilepsy; raised intra-ocular pressure including glaucoma; prostatic hypertrophy; severe hypertension or cardiovascular disease; bronchitis, bronchiectasis or asthma; hepatic impairment; renal impairment.
- Children and the elderly are more likely to experience the neurological anticholinergic effects and paradoxical excitation (eg. increased energy, restlessness, nervousness). Should not be used with other antihistamine containing products, including antihistamine containing cough and cold medicines.

4.5 Drug interactions

Phenylephrine: The co-administration of Monoamine Oxidase Inhibitors (MAOIs) or tricyclic antidepressants and an indirect or mixed-acting sympathomimetic may result in a hypertensive crisis and hence such concomitant use is best avoided. Additionally sympathomimetic may reduce the efficacy of beta-blocking and anti-hypertensive drugs.

Not recommended for patients currently receiving or within two weeks of stopping therapy with monoamine oxidase inhibitors.

Digoxin and cardiac glycosides: concomitant use of phenylephrine may increase the risk of irregular heartbeat or heart attack.

Chlorpheniramine: Concurrent use of Chlorpheniramine and hypnotics or anxiolytics may cause an increase in sedative effects; therefore, medical advice should be sought before taking Chlorpheniramine concurrently with these medicines. Chlorpheniramine inhibits phenytoin metabolism and can lead to phenytoin toxicity. The anti-cholinergic effects of Chlorpheniramine are intensified by MAOIs.

4.6 Use in special population

- Pediatric: Safe.
- Geriatric: Elderly population may be at greater risk for the side-effects.
- Liver impairment: Use with caution.
- Renal failure: Use with caution.

- Pregnancy and lactation: US Food and Drug Administration (FDA) has specified Chlorphenamine maleate as a pregnancy category B drug which indicates that animal and human studies have failed to demonstrate a risk to the fetus in any trimester. Sinarest AF Jr. syrup is recommended to be taken during pregnancy only under doctor's recommendation as safety of use of Sinarest AF Jr. syrup in pregnancy and lactation has not been established.

4.7Effects on ability to drive and use machine

It is advisable not to drive or operate machinery when on treatment with Sinarest AF Jr. syrup.

4.8Undesirable effects

Sinarest AF Jr. syrup is generally well tolerated and adverse events are rare.

Anaphylactic reaction, Thrombocytopenia, Palpitations, Tachycardia, Bradycardia, Tinnitus, Vertigo, Periorbital oedema, Vision blurred, Periorbital swelling, Eye swelling, Conjunctivitis, Nausea, Vomiting Diarrhoea, Dry mouth, Dyspepsia, Constipation, Abdominal pain, Hepatic function abnormal , Decreased appetite, Back pain, Myalgia, Dysuria, Urinary retention, Dyspnoea, Rash, Pruritus , Urticaria, Angioedema, Diabetes insipidus, Metabolic acidosis, Lactic acidosis, Headache, Dizziness.

4.G Overdose

There is limited experience of overdose with Sinarest AF Jr. syrup. Initiate general symptomatic and supportive measures in all cases of overdosages where necessary.

5 Pharmacological properties

5.1 Mechanism of action

Phenylephrine decreases nasal congestion by acting on α_1 -adrenergic receptors in the arterioles of the nasal mucosa to produce constriction; this leads to decreased edema and increased drainage of the sinus cavities.

In allergic reactions an allergen interacts with and cross-links surface IgE antibodies on mast cells and basophils. Once the mast cell-antibody-antigen complex is formed, a complex series of events occurs that eventually leads to cell-degranulation and the release of histamine (and other chemical mediators) from the mast cell or basophil. Once released, histamine can react with local or widespread tissues through histamine receptors. Histamine, acting on H_1 -receptors, produces pruritis, vasodilatation, hypotension, flushing, headache, tachycardia, and bronchoconstriction. Histamine also increases vascular permeability and potentiates pain. Chlorpheniramine maleate binds to the histamine H_1 receptor. This block the action of endogenous histamine, which subsequently leads to temporary relief of the negative symptoms brought on by histamine.

The Sodium citrate in Sinarest AF Jr. Syrup liquefies mucus and helps expectoration.

Menthol has a cooling and soothing effect.

5.2 Pharmacodynamic properties

Phenylephrine is a powerful vasoconstrictor. It is used as a nasal decongestant and cardiostimulant agent. Phenylephrine is a postsynaptic α_1 -receptor agonist with little effect on β -receptors of the heart. Parenteral administration of Phenylephrine causes a rise in systolic

and diastolic pressures, a slight decrease in cardiac output, and a considerable increase in peripheral resistance; most vascular beds are constricted, and renal, splanchnic, cutaneous, and limb blood flows are reduced while coronary blood flow is increased. Phenylephrine also causes pulmonary vessel constriction and subsequent increase in pulmonary arterial pressure. Vasoconstriction in the mucosa of the respiratory tract leads to decreased edema and increased drainage of sinus cavities.

Chlorpheniramine maleate is a histamine H_1 antagonist of the alkylamine class. It competes with histamine for the normal H_1 -receptor sites on effector cells of the gastrointestinal tract, blood vessels and respiratory tract. It provides effective, temporary relief of sneezing, watery and itchy eyes, and runny nose due to hay fever and other upper respiratory allergies.

5.3 Pharmacokinetic properties

Phenylephrine has low oral bioavailability owing to irregular absorption and first-pass metabolism by monoamine oxidase in the gut and liver. When injected subcutaneously or intramuscularly it takes 10 to 15 minutes to act; subcutaneous and intramuscular injections are effective for up to about 1 hour and up to about 2 hours, respectively. Intravenous injections are effective for about 20 minutes. Systemic absorption follows topical application.

Chlorphenamine maleate is absorbed relatively slowly from the gastrointestinal tract, peak plasma concentrations occurring about 2.5 to 6 hours after oral doses. Bioavailability is low, values of 25 to 50% having been reported. Chlorphenamine appears to undergo considerable first-pass metabolism. About 70% of Chlorphenamine in the circulation is bound to plasma proteins. There is wide inter individual variation in the pharmacokinetics of Chlorphenamine; values ranging from 2 to 43 hours have been reported for the half-life. Chlorphenamine is widely distributed in the body and enters the CNS. Chlorphenamine maleate is extensively metabolised. Metabolites include desmethylchlorpheniramine and didesmethylchlorphenamine. Unchanged drug and metabolites are excreted primarily in the urine; excretion is dependent on urinary pH and flow rate. Only trace amounts have been found in the faeces. Duration of action of 4 to 6 hours has been reported; this is shorter than may be predicted from pharmacokinetic parameters. More rapid and extensive absorption, faster clearance, and a shorter half-life have been reported in children.

6 Nonclinical properties


6.1 Animal Toxicology or Pharmacology

Not required.

7. Description


8. Pharmaceutical particulars

Phenylephrine is in a class of medications called nasal decongestants. Its chemical name is hydrogen 3-[(1*R*)-1-hydroxy-2-(methylamino)ethyl]phenol and its structural formula is:



Its empirical formula is $C_9H_{13}NO_2$, and its molecular weight is 167.20 g/mol.

Chlorpheniramine maleate is in a class of medications called antihistamines. Its chemical name is (Z)-but-2-enedioic acid;3-(4-chlorophenyl)-N,N-dimethyl-3-pyridin-2-ylpropan-1-amine and its structural formula is :



Its empirical formula is $C_{16}H_{19}ClN_2 \cdot C_4H_4O_4$ or $C_{20}H_{23}ClN_2O_4$ and its molecular weight is 390.9 g/mol

- 8.1 **Incompatibilities:** There are no known incompatibilities.
- 8.2 **Shelf-life:** 24 months
- 8.3 **Packaging Information:** Sinarest AF Jr. syrup is available in bottles of 75 ml.
- 8.4 **Storage and handling instructions:** Store below 25 °C in a dark place. Keep the bottle tightly closed.

9. Patient Counselling Information

9.1 **Adverse reactions:** Refer part 4.8

9.2Drug Interactions: Refer part 4.5

9.3Dosage: Refer part 4.2

9.4Storage: Refer part 8.4

9.5Risk factors: Refer part 4.4

9.6Self-monitoring information: NA

9.7Information on when to contact a health care provider or seek emergency help:

Patient is advised to be alert for the emergence or worsening of the adverse reactions and contact the prescribing physician.

9.8Contraindications: Refer part 4.3

10. Details of manufacturer:

CENTAUR PHARMACEUTICALS PVT. LTD.

Plant I, Plot No. 3, 5B, 2C, Tivim

Industrial Estate, Karaswada, Mapusa,

Goa- 403 526.

Regd. Off.: Centaur House,

Santacruz (E), Mumbai- 400 055.

11. License number date: 175

12. Date of revision: April 2025