

CatarestTM

Composition:

Potassium Iodide IP	3.3 % W/V
Sodium Chloride IP	0.83 % W/V
Calcium Chloride IP	1.0 % W/V

Indication:

- Catarest Eye Drops can be used for prevention of lenticular opacity pre-operatively and for prevention of Posterior Capsular Opacity (PCO), post-operatively, in patients with cataract. Especially, when the decision is to postpone to cataract surgery in patients with uncontrolled hypertension, diabetes mellitus, severe bleeding disorders and severe respiratory disorders.
- Catarest eye drops delays cataractogenesis, maintains normal nutrition of lens, assists visual improvement in early cataract and prevents posterior capsular opacification.

Contraindication:

Catarest is not advocated in those with known history of hypersensitivity to its ingredients

Drug Interaction:

Not any.

Adverse effects:

Catarest adverse effects are not reported.

Use in special population:

1. **Pediatric:** No data available.
2. **Geriatric:** No data available.
3. **Liver impairment:** No data available.
4. **Renal failure:** No data available.
5. **Pregnancy and lactation:** Doctor advice is recommended.

Dosage:

As directed by physician.

Presentation:

Catarest eye drops is available in 10ml lupolen vial.

Storage and handling:

Store in cool and dry place.



We Impart Health to Life