SINAREST NASAL SPRAY

Composition

Oxymetazoline 0.05% w/v

Sorbitol 2.0% w/v

Pharmacology

Oxymetazoline in SINAREST Nasal Spray is a long acting and topical nasal decongestant. It acts by stimulating the α-adrenergic receptors of the vascular smooth muscle. This leads to constriction of the dilated arteriolar network within the nasal mucosa and thus reduces blood flow in the engorged edematous nasal area. The constriction results in shrinkage of the engorged mucous membrane, which promotes drainage, improves nasal ventilation and relieves the feeling of stuffiness.

SINAREST Nasal Spray produces rapid, long lasting and predictable nasal decongestant effect.

Indications

SINAREST NASAL Spray is indicated for:

Relief of nasal congestion due to a cold, upper respiratory allergies or sinusitis.

Contraindications

The use of SINAREST Nasal Spray is contraindicated in patients with known hypersensitivity to its ingredient.

Precautions

SINAREST Nasal Spray should be administered with caution in patients with hypertension, coronary artery disease, hyperthyroidism or diabetes mellitus.

As with formulations, the use of the same packing of SINAREST Nasal Spray by more than one person may spread infection.

Pregnancy & Lactation

The safety of use of SINAREST Nasal Spray in pregnancy and lactation has not been established. Therefore, use only when clearly indicated.

Drug Interactions

Clinically significant drug interactions may occur on concomitant administration of SINAREST Nasal Spray with monoamine oxidase inhibitors, tricyclic antidepressants, b-adrenergic agents, methyldopa, reserpine and veratrum alkaloids.

Adverse Effects

SINAREST Nasal Spray may occasionally cause local stinging or burning sensation, sneezing, and dryness of the mouth and throat. Prolonged use may cause rebound congestion and drug induced rhinitis.

Dosage & Administration

The usual recommended dose of SINAREST Nasal Spray in adults is 2-3 sprays into each nostril bid.

Presentation

SINAREST Nasal Spray is available in 10 ml bottle.