

Sinarest[®] AF PLUS SYRUP

1. Composition

Phenylephrine	5 mg
Chlorpheniramine maleate	2mg
Sodium Citrate	60mg

2. Dosage form and strength

Sinarest AF Plus Syrup is available in 60 ml bottle.

3. Clinical particulars

3.1 Therapeutic indication

Sinarest AF Plus Syrup is indicated for cold without fever and allergic rhinitis.

3.2 Posology and method of administration

As directed by physician.

3.3 Contraindication

The use of Sinarest AF Plus Syrup is contraindicated in patients with:

- Hypersensitivity to any of the ingredients of the formulation.
- Severe hypertension.

3.4 Special warnings and precautions for use

- In case a hypersensitivity reaction occurs which is rare, Sinarest AF Plus Syrup should be discontinued.
- Sinarest AF Plus Syrup contains Paracetamol and therefore should not be used in conjunction with other Paracetamol containing products.
- Sinarest AF Plus Syrup should be used with caution in patients with renal or hepatic dysfunction, diabetes mellitus, hyperthyroidism, cardiovascular problems, epilepsy and closed angle glaucoma.

3.5 Drug interactions



We Impart Health to Life

Clinically significant drug interactions may occur on concomitant administration of Sinarest AF Plus Syrup with monoamine oxidase inhibitors, tricyclic antidepressants, beta-adrenergic agents, and methyldopa, reserpine and veratrum alkaloids.

3.6 Use in special population

- Pediatric: Safe.
- Geriatric: Elderly population may be at greater risk for the side-effects.
- Liver impairment: Use with caution.
- Renal failure: Use with caution.
- Pregnancy and lactation: US Food and Drug Administration (FDA) has specified Chlorpheniramine maleate as a pregnancy category B drug which indicates that animal and human studies have failed to demonstrate a risk to the fetus in any trimester. Paracetamol has been specified as a pregnancy category C drug which indicates that animal studies show an adverse effect on the fetus but there are no teratogenic studies of Paracetamol in pregnant women. Sinarest is recommended to be taken during pregnancy only under doctor's recommendation.

3.7 Effects on ability to drive and use machine

Chlorpheniramine in Sinarest AF Plus Syrup may cause sedation. It is advisable not to drive or operate machinery when on treatment with Sinarest AF Plus Syrup.

3.8 Undesirable effects

Sinarest AF Plus Syrup is generally well tolerated and adverse events are rare. Hypersensitive individuals may display ephedrine-like reactions such as tachycardia, palpitations, headache, dizziness and nausea. Use of sympathomimetics has been associated with fear, anxiety, restlessness, tremor, weakness, dysuria, insomnia, hallucinations and convulsions. Chlorpheniramine in Sinarest AF Plus Syrup may cause sedation.

3.9 Overdose

There is limited experience of overdose with Sinarest AF Plus Syrup. Initiate general symptomatic and supportive measures in all cases of overdosages where necessary.

4. Pharmacological properties

4.1 Mechanism of action

Phenylephrine decreases nasal congestion by acting on α 1-adrenergic receptors in the arterioles of the nasal mucosa to produce constriction; this leads to decreased edema and increased drainage of the sinus cavities.

In allergic reactions an allergen interacts with and cross-links surface IgE antibodies on mast cells and basophils. Once the mast cell-antibody-antigen complex is formed, a complex series of events occurs that eventually leads to cell-degranulation and the release of histamine (and other chemical mediators) from the mast cell or basophil. Once released, histamine can react with local or widespread tissues through histamine receptors. Histamine, acting on H1-receptors, produces pruritis, vasodilatation, hypotension, flushing, headache, tachycardia, and bronchoconstriction. Histamine also increases vascular permeability and potentiates pain. Chlorpheniramine maleate binds to the histamine H1 receptor. This block the action of endogenous histamine, which subsequently leads to temporary relief of the negative symptoms brought on by histamine.

The Sodium citrate in Sinarest AF Plus Syrup liquefies mucus and helps expectoration.

4.2 Pharmacodynamic properties

Phenylephrine is a powerful vasoconstrictor. It is used as a nasal decongestant and cardiostimulant agent. Phenylephrine is a postsynaptic α 1-receptor agonist with little effect on β -receptors of the heart. Parenteral administration of phenylephrine causes a rise in systolic and diastolic pressures, a slight decrease in cardiac output, and a considerable increase in peripheral resistance; most vascular beds are constricted, and renal, splanchnic, cutaneous, and limb blood flows are reduced while coronary blood flow is increased. Phenylephrine also causes pulmonary vessel constriction and subsequent increase in pulmonary arterial pressure. Vasoconstriction in the mucosa of the respiratory tract leads to decreased edema and increased drainage of sinus cavities.

Chlorpheniramine maleate is a histamine H1 antagonist of the alkylamine class. It competes with histamine for the normal H1-receptor sites on effector cells of the gastrointestinal tract, blood vessels and respiratory tract. It provides effective, temporary relief of sneezing, watery and itchy eyes, and runny nose due to hay fever and other upper respiratory allergies.

Citrate prevents activation of the clotting cascade by chelating calcium ions. Citrate neutralizes acid in the stomach and urine, raising the pH.

4.3 Pharmacokinetic properties

Phenylephrine has low oral bioavailability owing to irregular absorption and first-pass metabolism by monoamine oxidase in the gut and liver. When injected subcutaneously or intramuscularly it takes 10 to 15 minutes to act; subcutaneous and intramuscular injections are effective for up to about 1 hour and up to about 2 hours, respectively. Intravenous

injections are effective for about 20 minutes. Systemic absorption follows topical application.

Chlorphenamine maleate is absorbed relatively slowly from the gastrointestinal tract, peak plasma concentrations occurring about 2.5 to 6 hours after oral doses. Bioavailability is low, values of 25 to 50% having been reported. Chlorphenamine appears to undergo considerable first-pass metabolism. About 70% of chlorphenamine in the circulation is bound to plasma proteins. There is wide inter individual variation in the pharmacokinetics of chlorphenamine; values ranging from 2 to 43 hours have been reported for the half-life. Chlorphenamine is widely distributed in the body, and enters the CNS. Chlorphenamine maleate is extensively metabolised. Metabolites include desmethyl- and didesmethylchlorphenamine. Unchanged drug and metabolites are excreted primarily in the urine; excretion is dependent on urinary pH and flow rate. Only trace amounts have been found in the faeces. Duration of action of 4 to 6 hours has been reported; this is shorter than may be predicted from pharmacokinetic parameters. More rapid and extensive absorption, faster clearance, and a shorter half-life have been reported in children.

Sodium citrate is metabolised after absorption to bicarbonate. Bicarbonate ions are excreted in the urine, which is rendered alkaline, and there is an accompanying diuresis.

5. Nonclinical properties

5.1 Animal Toxicology or Pharmacology

Not required.

6. Description

Already mentioned and covered in the above points.

7. Pharmaceutical particulars

7.1 Incompatibilities

There are no known incompatibilities.

7.2 Shelf-life

24 months.

7.3 Storage and handling instructions

Store below 30 °C in a dark and dry place.