

Zincogut[™]

Syrup

1. Composition

Each 5ml contains

Zinc Gluconate equivalent to elemental Zinc 20mg

2. Dosage form and strength

Zincogut is presented as 60ml Bottle.

3. Clinical particulars

3.1 Therapeutic indication

Zincogut is indicated in patients with:

- Acute Infective Diarrhoea
- Persistent Diarrhoea

3.2 Posology and method of administration

- For Infants between 2-6 months =2.5 ml for 14 days
- For Children between 6months-5years = 5 ml for 14 days
- To be taken once a day

3.3 Contraindication

Zincogut is contraindicated in patients:

- Hypersensitive to components of the formulation.
- Infected with Human Immuno Deficiency Virus (HIV)

3.4 Special warnings and precautions for use

- Infants and children: Zinc is likely safe when taken by mouth appropriately in the recommended amounts.
- Diabetes: Large doses of zinc can lower blood sugar in people with diabetes.

3.5 Drug interactions

- Antibiotics (Quinolone antibiotics) interacts with ZINC: Zinc might decrease how much antibiotic the body absorbs. Taking zinc along with some antibiotics might decrease the effectiveness of some antibiotics. To avoid this interaction take zinc supplements at least 1 hour after antibiotics.



We Impart Health to Life

- Antibiotics (Tetracycline antibiotics) interacts with ZINC: Zinc can attach to tetracycline in the stomach. This decreases the amount of tetracycline that can be absorbed. Taking zinc with tetracycline might decrease the effectiveness of tetracycline.
- Cisplatin (Platinol-AQ) interacts with ZINC
- Penicillamine interacts with ZINC.

3.6 Use in special population

- Pediatric: Safe in children.
- Geriatric: Problems in older adults have not been reported with intake of normal daily recommended amounts.
- Liver impairment: No data available.
- Renal failure: No data available.
- Pregnancy and lactation: Although there appears to be no harmful effects of zinc supplementation in pregnancy and lactation, consultation with doctor is recommended.

3.7 Effects on ability to drive and use machine

No data available.

3.8 Undesirable effects

- Metallic taste
- Abdominal pain
- Diarrhoea
- Vomiting
- Interferes with iron absorption causing Anaemia.
- Zinc can interfere with Copper absorption

3.9 Overdose

There is limited experience of overdose with Zincogut syrup. Initiate general symptomatic and supportive measures in all cases of overdosages where necessary.

4. Pharmacological properties

4.1 Mechanism of action

Zinc helps in diarrhoea by causing:

- ✓ Improved absorption of water and electrolytes by the intestine and increased levels of enterocyte brush border enzymes.
- ✓ Zinc reduces stool volume and duration of diarrhoea.



We Impart Health to Life

- ✓ Action on intestinal mucosa: Zinc reduces fluid secretion in the intestine by indirectly inhibiting cAMP dependent Cl transport across the mucosa through an action on the basolateral membrane K⁺.
- ✓ In terms of Immune Response: Zinc strengthens the immune response and helps in regeneration of intestinal epithelium.

4.2 Pharmacodynamics properties

- Zinc is an important mineral found in almost every cell in the human body. It promotes the activity of about 100 enzymes. Zinc deficiency is often associated with an increased risk of infection. When they are used to treat the common cold, zinc supplements may interfere with rhinovirus cleavage or adhesion and may play a role in protecting plasma membranes from microbial toxins and complement.

4.3 Pharmacokinetic properties

- Zinc is absorbed in the small intestine by a carrier-mediated mechanism. More recent studies have determined different absorption rates for various populations based on their type of diet and phytate to zinc molar ratio. Zinc-deprived humans absorb this element with increased efficiency, whereas humans on a high-zinc diet show a reduced efficiency of absorption. Approximately 60-70% of the zinc in circulation is bound to albumin. Any condition that alters serum albumin concentration may have a secondary effect on serum zinc levels. Considerable amounts of zinc are secreted through both biliary and intestinal secretions, however most is reabsorbed.

5. Nonclinical properties

5.1 Animal Toxicology or Pharmacology

Not required.

6. Description

Already mentioned and covered in the above points.

7. Pharmaceutical particulars

7.1 Incompatibilities

There are no known incompatibilities.

7.2 Shelf-life

24 months

7.3 Storage and handling instructions



We Impart Health to Life

Store at 25°C in cool and dry place.



We Impart Health to Life